



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2015

---

## **No association between the ALDH2 promoter polymorphism rs886205, alcohol dependence, and risky alcohol consumption in a German population**

Nassab, Mani Haschemi ; Rhein, Mathias ; Heese, Peter ; Glahn, Alexander ; Frieling, Helge ; Linnebank, Michael ; Bleich, Stefan ; Kornhuber, Johannes ; Heberlein, Annemarie ; Grallert, Harald ; Peters, Annette ; Rawal, Rajesh ; Strauch, Konstantin ; Hillemacher, Thomas

DOI: <https://doi.org/10.1097/YPG.0000000000000073>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-101285>

Journal Article

Published Version

Originally published at:

Nassab, Mani Haschemi; Rhein, Mathias; Heese, Peter; Glahn, Alexander; Frieling, Helge; Linnebank, Michael; Bleich, Stefan; Kornhuber, Johannes; Heberlein, Annemarie; Grallert, Harald; Peters, Annette; Rawal, Rajesh; Strauch, Konstantin; Hillemacher, Thomas (2015). No association between the ALDH2 promoter polymorphism rs886205, alcohol dependence, and risky alcohol consumption in a German population. *Psychiatric Genetics*, 25(1):41-42.

DOI: <https://doi.org/10.1097/YPG.0000000000000073>

# No association between the ALDH2 promoter polymorphism rs886205, alcohol dependence, and risky alcohol consumption in a German population

Mani Haschemi Nassab<sup>a,b</sup>, Mathias Rhein<sup>b</sup>, Peter Heese<sup>c</sup>, Alexander Glahn<sup>a</sup>, Helge Frieling<sup>a,b</sup>, Michael Linnebank<sup>d</sup>, Stefan Bleich<sup>a</sup>, Johannes Kornhuber<sup>e</sup>, Annemarie Heberlein<sup>a</sup>, Harald Grallert<sup>f,g</sup>, Annette Peters<sup>g</sup>, Rajesh Rawal<sup>h</sup>, Konstantin Strauch<sup>h,i</sup> and Thomas Hillemacher<sup>a</sup>

Psychiatric Genetics 2015, 25:41–42

<sup>a</sup>Department of Psychiatry, Socialpsychiatry and Psychotherapy, Center for Addiction Research (CARE), <sup>b</sup>Molecular Neurosciences Laboratory, Department of Psychiatry, Socialpsychiatry and Psychotherapy, Hannover Medical School, Hannover, <sup>c</sup>Department of Addiction and Psychotherapy, LVR-Clinic Bonn, Bonn, <sup>d</sup>Department of Psychiatry and Psychotherapy, University Hospital, Friedrich-Alexander University of Erlangen-Nuremberg, Erlangen, <sup>e</sup>Research Unit of Molecular Epidemiology, <sup>f</sup>Institute of Genetic Epidemiology, Helmholtz Center Munich, German Research Center for Environmental Health, <sup>g</sup>Institute of Epidemiology II, Helmholtz Center Munich, German Center for Diabetes Research, Neuherberg, <sup>h</sup>Department of Genetic Epidemiology, Institute of Medical

Informatics, Biometry and Epidemiology, Ludwig-Maximilians Universität, Munich, Germany and <sup>i</sup>Department of Neurology, University Hospital Zürich, Zürich, Switzerland

Correspondence to Mani Haschemi Nassab, PhD, Molecular Neurosciences Laboratory, Department of Psychiatry, Socialpsychiatry and Psychotherapy, Hannover Medical School, Feodor-Lynen Straße 35, 30625 Hannover, Germany Tel: +49 511 532 7245; fax: +49 511 532 7276; e-mail: hascheminassab.mani@mh-hannover.de

Received 17 July 2014 Accepted 24 October 2014

Susceptibility to the negative effects of alcohol consumption, for example, headache, nausea, and flushing is associated with blood levels of toxic acetaldehyde, which is mainly eliminated by active aldehyde dehydrogenase 2 (ALDH2). A polymorphism in the coding region of the *ALDH2* gene, rs671, causes loss of enzymatic activity and protection against alcohol dependence, but is predominantly present in East-Asian populations (Brennan *et al.*, 2004). In contrast to rs671, the noncoding *ALDH2* promoter polymorphism rs886205 (A > G) appears in relevant frequency in different populations, including European, as a risk marker for alcohol-related carcinoma (Hashibe *et al.*, 2006), and is known to reduce *ALDH2* gene transcription and promoter activity *in vivo* and *in vitro* (Kimura *et al.*, 2009).

In a previous longitudinal study with 82 alcohol-dependent patients and 34 controls of German descent, we detected different rs886205 alleles and genotype frequencies between the groups, but not reaching significance (allele frequency:  $\chi^2 = 3.18$ ;  $P = 0.074$  and genotype frequency:  $\chi^2 = 2.89$ ;  $P = 0.089$ ). We calculated that replication of this genetic effect in a larger cohort of at least 300 patients and controls would have sufficient power to confirm a potential impact of rs886205 on the associated risk for alcohol dependence ( $\alpha = 0.05$ ;  $1 - \beta = 0.83$ ).

Therefore, we genotyped marker rs886205 in 352 alcohol-dependent patients according to ICD-10 (Heese *et al.*, 2012) and two independent control cohorts that included 2742 (KORA S3) and 3175 (KORA S4) population-based controls. All individuals were of German descent and provided written informed consent. Genotype frequencies were in Hardy–Weinberg equilibrium and were as follows (patients/control KORA

S3/control KORA S4): A/A = 66.5/68.9/68.5, A/G = 31.8/28.0/28.2, G/G = 1.7/3.3/3.1.

Neither genotype nor allele frequencies showed significant differences between patients and KORA S3 controls (genotype:  $\chi^2 = 0.82$ ,  $P = 0.365$ ; allele type:  $\chi^2 = 0.11$ ;  $P = 0.745$ ) and patients and KORA S4 controls (genotype:  $\chi^2 = 0.60$ ,  $P = 0.438$ ; allele type:  $\chi^2 = 0.02$ ;  $P = 0.888$ ).

To examine whether the rs886205 genotype might affect alcohol consumption (grams of ethanol per day) in alcohol-dependent patients and controls, we applied a logistic regression model using risky alcohol consumption as the dependent variable (male individuals > 30 g/day, female individuals > 20 g/day) and rs886205 genotype, age, and sex as independent variables. Although age and sex had a significant impact ( $P < 0.0001$ ), the rs886205 genotype did not affect risky alcohol consumption in patients and controls ( $P = 0.965$ ).

So far, an association between rs886205 and alcohol dependence was only investigated in East-Asian populations, with contradictory outcome, and biasing effects of strong linkage disequilibrium with rs671 (Harada *et al.*, 1999). It was also suggested that the role of the *ALDH2* promoter polymorphism might vary across populations because of large differences in allelic frequencies between East-Asian and European populations (Kimura *et al.*, 2006).

Our findings show for the first time that the functional *ALDH2* promoter polymorphism rs886205 does not affect risk for alcohol dependence and risky alcohol consumption in German populations.

## Acknowledgements

### Conflicts of interest

There are no conflicts of interest.

## References

- Brennan P, Lewis S, Hashibe M, Bell DA, Boffetta P, Bouchardy C, *et al.* (2004). Pooled analysis of alcohol dehydrogenase genotypes and head and neck cancer: a HuGE review. *Am J Epidemiol* **159**:1–16.
- Harada S, Okubo T, Nakamura T, Fujii C, Nomura F, Higuchi S, Tsutsumi M (1999). A novel polymorphism (–357 G/A) of the ALDH2 gene: linkage disequilibrium and an association with alcoholism. *Alcohol Clin Exp Res* **23**:958–962.
- Hashibe M, Boffetta P, Zaridze D, Shangina O, Szeszenia-Dabrowska N, Mates D, *et al.* (2006). Evidence for an important role of alcohol – and aldehyde-metabolizing genes in cancers of the upper aerodigestive tract. *Cancer Epidemiol Biomarkers Prev* **15**:696–703.
- Heese P, Linnebank M, Semmler A, Muschler MA, Heberlein A, Frieling H, *et al.* (2012). Alterations of homocysteine serum levels during alcohol withdrawal are influenced by folate and riboflavin: results from the German Investigation on Neurobiology in Alcoholism (GINA). *Alcohol Alcohol* **47**:497–500.
- Kimura M, Kimura S, Matsushita S, Kashima H, Higuchi S (2006). ALDH2 promoter polymorphism has no effect on the risk for alcoholism. *Alcohol Alcohol* **41**:368–371.
- Kimura Y, Nishimura FT, Abe S, Fukunaga T, Tanii H, Saijoh K (2009). A promoter polymorphism in the ALDH2 gene affects its basal and acetaldehyde/ethanol-induced gene expression in human peripheral blood leukocytes and HepG2 cells. *Alcohol Alcohol* **44**:261–266.